

Accidental Bowel Leakage

Frequently Asked Questions

Overview

What is accidental bowel leakage?

Accidental bowel leakage (ABL) is the loss of normal control of your [bowels](#) . It also is called [fecal incontinence](#) or anal incontinence. This condition leads to leakage of solid stool, liquid stool, mucus, or gas.

Accidental bowel leakage can have a significant effect on your quality of life. But there is hope for improvement with the right treatment.

What is normal bowel function?

Normal bowel function requires healthy muscles and nerves in the [rectum](#) and [pelvis](#) . The [large intestine](#) (also called the [colon](#)) must be able to form and store the stool until you can get to the bathroom. The [sphincter muscles](#) of the [anus](#) must be strong enough to stay closed until you need to pass stool or gas.

What can I do if my bowel function doesn't seem normal?

Talking about ABL may feel embarrassing, but telling your [obstetrician–gynecologist \(ob-gyn\)](#) is the first step in getting treatment. Many types of treatment are available depending on the cause of the problem.

Symptoms and Causes

What are the symptoms of accidental bowel leakage?

Symptoms of a bowel control problem may include:

- Leaking stool, mucus, or gas
- Strong or urgent need to have bowel movements
- Diarrhea
- Constipation
- Streaks or stains of stool on your underwear

Can digestive disorders cause accidental bowel leakage?

ABL can be caused by digestive disorders, including:

- Irritable bowel syndrome (IBS)
- Inflammatory bowel disease (IBD)
- Other chronic diseases

Can childbirth be a cause of bowel control problems?

In some women, bowel control problems often are the result of vaginal childbirth. As the [fetus](#) passes through the [vagina](#), the muscles and tissues of the rectum may be stretched or torn. The anal sphincter also can be injured during vaginal delivery.

Some women have short-term bowel control problems right after childbirth that improve within a few days or weeks. Other women do not develop a problem until later in life. Also, as a person ages, the anal sphincter may weaken. A minor problem in a younger woman can become worse with age.

What are the risk factors for developing accidental bowel leakage?

Some other common factors that may put you at high risk of ABL include:

- Chronic constipation
- Diarrhea
- Increased number of bowel movements (more than 21 stools a week)
- Often feeling an urgent need to have a bowel movement
- Not being able to control urine
- Obesity
- Smoking
- Older age
- Decreased physical activity
- Anal intercourse
- Other health conditions, such as [depression](#) , [diabetes mellitus](#) , and IBD
- Past [radiation therapy](#) to the anal region

Diagnosis

How is the cause of accidental bowel leakage found?

Finding the cause of ABL is an important step in finding the right treatment. Talking honestly with your ob-gyn offers the best chance of finding the right treatment for you.

Your ob-gyn may examine your vagina, anus, and rectum. This is done to look for signs of problems, such as loss of nerve reflexes or muscle tone. In some cases, tests may be needed.

How can I prepare for an office visit?

At your office visit, your ob-gyn should ask about your medical history and symptoms. You may be asked to keep a record each time you pass stool. Write down whether you had regular bowel movement or leakage of stool or gas. Also, write down any other symptoms and what you were doing at the time. This symptom record will help your ob-gyn diagnose why you are having problems.

You also should tell your ob-gyn about any prescription or over-the-counter products or herbal remedies you may be taking. Some medications and supplements can cause

constipation or diarrhea and may contribute to your symptoms.

You and your ob-gyn should review the results of your physical exam and any tests that might have been ordered. You also should talk about your symptom record. Describe your symptoms clearly to help your ob-gyn find the best treatment for you.

What tests are available for accidental bowel leakage?

The following are some of the tests that may be done to see how well the muscles in your anus and rectum are working:

- Anorectal manometry—This test uses a small sensing device to test the strength of the anal sphincter. The device is placed in the anus and records changes in pressure as you relax and tighten the muscles.
- Defecography—This X-ray test is used to study the rectum and anal canal during a bowel movement.
- Nerve tests—These tests check if the nerves to the rectum and anus are working well.
- Imaging tests, such as an [ultrasound exam](#)
- Tests to look inside your anus, rectum, and colon, including [colonoscopy](#)
- Tests of blood, stool, and urine

Treatment Options

How is accidental bowel leakage treated?

The type of treatment depends on the cause of the problem and how severe it is. Your ob-gyn may care for you, or you may be referred to a doctor who specializes in treating ABL, such as a gastroenterologist.

What can I do to help treat accidental bowel leakage?

Many women with ABL can improve symptoms with self-care. Changes in diet, taking medication, or doing special exercises can make a difference for many women.

Also, your doctor can give you advice about caring for the skin around the anus, which can become irritated. This irritation can lead to pain or itching. You can manage anal discomfort by

- avoiding foods that irritate the perianal skin
- using zinc oxide ointment around the anus
- wearing disposable underwear or using pads in your own underwear

Talk with your ob-gyn about approaches that are best for you. Report any skin breakdown, rashes, irritation, or itching.

How can a food diary help?

If you leak gas or have diarrhea, keeping a food diary can help you identify foods that cause these problems. If you find a pattern, you can avoid these foods. Your food diary should note what you eat, how much, and when. You also will keep track of your symptoms, what type of bowel movements you have, and when they occur.

Which foods can make accidental bowel leakage worse?

Dairy products and foods that contain gluten, a protein found naturally in wheat, rye, and barley, can cause gas and diarrhea in some people. Some artificial sweeteners also can cause diarrhea. Some people have trouble absorbing fatty foods. Other people have trouble with spicy foods, alcohol, and caffeine. Keeping a food diary can help you figure out which foods affect you.

Which foods can help with constipation?

If you have constipation, your ob-gyn may suggest you eat more fruits and vegetables, drink plenty of water, and add fiber. Fiber is found in plant foods. It is the part of the plant that your body cannot digest. Fiber can help prevent constipation by adding bulk to the stool, making it easier to pass.

How much fiber should I eat each day?

Adult women should eat about 25 grams of fiber daily. Foods that are high in fiber include

- the outer shell of whole grains like oats, corn, and wheat
- beans
- berries
- green, leafy vegetables

You also can add fiber by using fiber supplements.

Is medication an option?

In some cases, yes. You and your ob-gyn may choose a medication that fits your situation and that can best control your bowel leakage. Some over-the-counter medications can help relieve symptoms. Stronger medications that treat diarrhea, control gas leakage, or prevent constipation may be prescribed.

Can exercise help my symptoms?

Yes. [Kegel exercises](#) strengthen the muscles that surround the openings of the anus, [urethra](#) , and vagina. Doing these exercises regularly may improve incontinence. Your ob-gyn or a physical therapist can help you be sure you are doing these exercises correctly.

How are Kegel exercises done?

Kegel exercises help tone the muscles of the pelvic floor. These muscles support your urethra, [bladder](#) , [uterus](#) , and rectum. Strengthening these muscles may help improve bladder control. After delivery, exercises may help tighten vaginal muscles that are stretched from childbirth. Here is how Kegel exercises are done:

- Squeeze the muscles that you use to stop the flow of urine. This pulls the vagina and rectum up and back.
- Hold for 3 seconds, then relax for 3 seconds.
- Do 10 contractions three times a day.
- Increase your hold by 1 second each week. Work your way up to 10-second holds.

Make sure you are not squeezing your stomach, thigh, or buttock muscles. You also should breathe normally. Do not hold your breath as you do these exercises.

You can do Kegel exercises anywhere—while working, driving in your car, or watching television. But you should not do these exercises when you are urinating.

What is bowel training?

Bowel training involves training yourself to have regular bowel movements at the same time each day. Several techniques can be used. Your ob-gyn can tell you which technique is best for you based on your symptoms.

What is biofeedback training?

When used to manage ABL, [biofeedback](#) is a training technique that helps you improve the function of the anal sphincter muscles. In biofeedback, sensors are placed inside or outside the anus. These sensors provide visual feedback on a monitor so you can see if you are contracting and relaxing muscles correctly.

Like exercising any other type of muscle, you can strengthen the anal sphincter muscles over time. By helping you identify your anal muscles, biofeedback can improve your ability to sense stool or gas in the rectum.

What devices can be used to manage symptoms?

- One device pumps air into a balloon inside your vagina. The inflated balloon puts pressure on the rectum. This keeps stools from passing. To control when you pass stool, you can add or remove air from the balloon.
- Another device is an anal plug. You can remove the plug when you need to pass stool.

What is sacral nerve stimulation?

Electrical stimulation therapy can be used when the nerves that control the bowels are not working correctly. With this treatment, a device is implanted under the skin, usually right above the buttocks. A thin wire is placed near the sacral nerves (near the tailbone), which control the colon, rectum, and anal sphincter. The device sends a mild electrical signal along the wire to these nerves to restore the normal function of the bowels.

How are injections used to treat accidental bowel leakage?

A special kind of injection can be used to treat bowel control problems. With this treatment, an agent that adds bulk is injected in the tissue surrounding the anus. This narrows the anal opening to help the anal sphincter work better.

Can surgery treat accidental bowel leakage?

In some cases, surgery may help correct loss of bowel control. Most people will try other treatments before surgery is recommended.

If the sphincter muscle of the anus is torn, it sometimes can be repaired with surgery. Sutures (stitches) are used to bring together the torn ends of the sphincter so they can heal. Your ob-gyn can help you decide if surgery is an option for you.

Glossary

Anus: The opening of the digestive tract through which bowel movements leave the body.

Biofeedback: A technique used by physical therapists to help a person control body functions, such as heartbeat or blood pressure.

Bladder: A hollow, muscular organ in which urine is stored.

Bowels: The small and large intestines.

Colon: The large intestine.

Colonoscopy: An exam of the large intestine using a small, lighted instrument.

Depression: Feelings of sadness for periods of at least 2 weeks.

Diabetes Mellitus: A condition in which the levels of sugar in the blood are too high.

Fecal Incontinence: Involuntary loss of control of the bowels.

Fetus: The stage of human development beyond 8 completed weeks after fertilization.

Inflammatory Bowel Disease (IBD): The name for a group of diseases that cause inflammation of the intestines. Examples include Crohn's disease and ulcerative colitis.

Irritable Bowel Syndrome (IBS): A digestive disorder that can cause gas, diarrhea, constipation, and belly pain.

Kegel Exercises: Pelvic muscle exercises. Doing these exercises helps with bladder and bowel control as well as sexual function.

Large Intestine: The portion of the intestine that starts at the end of the small intestine and ends at the anus.

Obstetrician–Gynecologist (Ob-Gyn): A doctor with special training and education in women's health.

Pelvis: The lower portion of the trunk of the body.

Radiation Therapy: Treatment with radiation.

Rectum: The last part of the digestive tract.

Sphincter Muscles: Muscles that can close a bodily opening, such as the sphincter muscle of the anus.

Ultrasound Exam: A test in which sound waves are used to examine inner parts of the body.

Urethra: A tube-like structure. Urine flows through this tube when it leaves the body.

Uterus: A muscular organ in the female pelvis. During pregnancy, this organ holds and nourishes the fetus. Also called the womb.

Vagina: A tube-like structure surrounded by muscles. The vagina leads from the uterus to the outside of the body.

If you have further questions, contact your ob-gyn.

Don't have an ob-gyn? [Learn how to find a doctor near you.](#)

FAQ139

Last updated: November 2021

Last reviewed: April 2023

Copyright 2024 by the American College of Obstetricians and Gynecologists. All rights reserved. Read [copyright and permissions information](#).

This information is designed as an educational aid for the public. It offers current information and opinions related to women's health. It is not intended as a statement of the standard of care. It does not explain all of the proper treatments or methods of care. It is not a substitute for the advice of a physician. Read [ACOG's complete disclaimer](#).

About ACOG

Disclaimer

Contact Us

How to Find an Ob-Gyn



Copyright 2024 American College of Obstetricians and Gynecologists

Privacy Statement

|

Terms and Conditions of Use